

CARDIOSPERMUM CORINDUM L. (SAPINDACEAE) HAS GASTROPROTECTIVE AND ANTISPASMODIC EFFECTS IN RODENT MODEL

CARDIOSPERMUM CORINDUM L. (SAPINDACEAE) TEM EFEITO PROTETOR GÁSTRICO E ANTIESPASMÓDICO EM MODELO DE ROEDORES

Joelmir Lucena Veiga da Silva¹, Valdelena Alessandra Silva², Pamela França Nascimento³, Dulce de Fátima Ribeiro Suzart³, Fabiana Lima Silva^{4,5}, José Maria Barbosa-Filho⁶, Vera Lúcia Silva Rigoni⁷, Viviane Louise Andree Nouailhetas⁸

¹ Faculdade de Medicina de Olinda, Olinda, Pernambuco, Brazil. ² Universidade Nove de Julho, São Paulo, São Paulo, Brazil. ³ Universidade Nove de Julho, São Paulo, São Paulo, Brazil. ⁴ Instituto de Química, Universidade de São Paulo, São Paulo, São Paulo, Brazil. ⁵ Instituto de Ciências da Saúde, Universidade Paulista, São Paulo, São Paulo, Brazil. ⁶ Departamento de Ciências Farmacêuticas, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brazil. ⁷ Departamento de Biofísica, Universidade Federal de São Paulo, São Paulo, São Paulo, Brazil. ⁸ Departamento de Biofísica, Universidade Federal de São Paulo, São Paulo, São Paulo, Brazil.

ABSTRACT

Objectives: To investigate the effects of the hydroethanolic extract of aerial parts of *Cardiospermum corindum* (Cc-EtOH) in the ethanol-induced gastric ulcer assay in rats, its oral toxicity in mice, and its antispasmodic activity in isolated rat ileum *in vitro*. **Methods:** The anti-ulcerative activity of Cc-EtOH was evaluated in ethanol-induced rat gastric lesions. Oral toxicity was assessed in mice following a single acute treatment and monitored for 14 days. The antiespasmotic effect of Cc-EtOH in isolated rat ileum was analyzed *in vitro*. **Results:** Ethanol-induced gastric lesions in rats (ulcer index = 367.5 ± 89.3 mm²) were prevented by 50, 150, and 500 mg/kg Cc-EtOH (ulcer index = 210.0 ± 55.8 , 119.2 ± 39.3 , and 47.7 ± 13.2 mm², respectively), with efficacy comparable to omeprazole. Histological findings corroborated these results. A single oral dose of 2 g/kg Cc-EtOH induced transient sedation in mice after 60 min (n = 5), which was reversed within 30 min. No acute toxicity was observed, as evidenced by the absence of animal mortality, changes in body weight, or alterations in water or food intake over 14 days. Cc-EtOH reduced *in vitro* phasic contractions of isolated rat ileum elicited by carbachol or KCl (IC₅₀ = 236 ± 59 and 201 ± 43 mg/mL, respectively). **Conclusion:** The aerial parts of *C. corindum* contain bioactive compounds that exert effective gastric protection in rats, show no evidence of acute toxicity, and display antispasmodic effect *in vitro*.

Keywords: Medicinal plant; Plant extract; Anti-ulcer Agent; Antispasmodic effect; Toxicity

RESUMO

Objetivos: Investigar os efeitos do extrato de etanol bruto obtido das partes aéreas *C. corindum* (Cc-EtOH) sobre o teste de úlcera gástrica induzido pelo etanol em ratos, sua toxicidade aguda oral em camundongos e sua atividade antiespasmódica em íleo isolado de rato. **Métodos:** O Cc-EtOH foi avaliado em lesões gástricas induzidas por etanol. Para verificar a segurança deste extrato *in vivo*, o teste de toxicidade aguda foi realizado durante 14 dias em camundongos. Finalmente, os efeitos do extrato no íleo do rato foram analisados. **Resultados:** O extrato Cc-EtOH (50, 150 e 500 mg/kg) protegeu significativamente a camada de mucosa gástrica do rato (área da úlcera = $210,0 \pm 55,8$; $119,2 \pm 39,3$ e $47,7 \pm 13,2$ mm², respectivamente) das lesões induzidas pelo etanol (área da úlcera = $367,5 \pm 89,3$ mm²), tão efetivamente como omeprazol. As análises histológicas também corroboraram seu efeito protetor. Além disso, tratamento de 60 minutos com uma dose única de 2g/kg v.o., o extrato de Cc-EtOH promoveu a sedação dos camundongos (n = 5), que se reverteu após 30 minutos, sem causar nem a morte do animal, nem a massa corporal, a ingestão de água, nem as alterações da alimentação, durante 14 dias, mostrando não toxicidade aguda. O efeito antiespasmódico do extrato Cc-EtOH foi evidenciado no componente fásico das contrações induzidas por CCh e KCl (IC₅₀ = 236 ± 59 e 201 ± 43 mg/mL, respectivamente) em íleo isolado do rato. **Conclusão:** As partes aéreas do *C. corindum* contêm compostos químicos que são capazes de promover proteção gástrica efetiva em ratos, na ausência de qualquer toxicidade aguda, combinada com o efeito antiespasmódico *in vitro*.

Palavras-chave: Planta medicinal; Extrato vegetal; Agente anti-úlcera; Efeito antiespasmódico; Toxicidade

INTRODUCTION

Gastric hyperacidity and gastroduodenal ulcers are frequent conditions that currently represent a major worldwide public health concern¹. *Cardiospermum corindum* L. (*Sapindaceae*), popularly known as *balãozinho*, occurs in the northeastern, southeastern, and southern regions of Brazil². In traditional medicine, particularly in the Brazilian northeast, infusions prepared from *C. corindum* are used to treat liver disorders and rheumatism, as a memory tonic, diuretic, and emmenagogue³. Several compounds have been isolated and chemically identified from the leaves and other aerial parts of *C. corindum*^{4,5}. Despite its widespread folk use for stomach ailments, the biological activity of its compounds has not yet been scientifically described.

The present study aimed to investigate the effects of the hydroethanolic extract obtained from the aerial parts of *C. corindum* on ethanol-induced gastric ulcers in rats, its acute toxicity in mice, and its antispasmodic activity in isolated rat ileum.

METHODS

The aerial parts of *Cardiospermum corindum* L. were collected during the flowering period at the base of Pico do Jabre, Paraíba, Brazil. The plant was identified, and a voucher specimen (No. M.F. Agra et al. 6898) was deposited in the Prof. Lauro Pires Xavier herbarium at Universidade Federal da Paraíba. The crude hydroethanolic extract (Cc-EtOH) was prepared according to previous report⁵.

The Cc-EtOH extract was dissolved in Tween-20 (0.32 mg/mL) and diluted in distilled water to final doses. Wistar rats (200 - 300 g) were stratified into six groups (n = 5) and pretreated with 50, 150, 250, or 500 mg/kg Cc-EtOH extract (gavage), 4 mg/kg omeprazole (intraperitoneal), or vehicle (distilled water-Tween-20, 10 mL/kg). One hour later, 1 mL of absolute ethanol was orally administered to each animal. After one hour, animals were euthanized in a CO₂ chamber, and their stomachs were collected and dissected along the greater curvature⁶. The ulcer index (UI) was assessed by quantifying the extent of erosion and experimental gastric lesions⁷. Samples were processed for histological analysis and stained with hematoxylin and eosin. This protocol was approved by the ethics committee on animal use of Universidade Nove de Julho (opinion AN 0002/11).

Swiss male mice (25 - 30 g) were divided into two groups (n = 5) and orally treated with either a single dose of 2 g/kg Cc-EtOH extract ("treated-group") or vehicle (10 mL/kg distilled water-Tween-20; "control group"). Animals were observed for 120 min at 30 min intervals to assess toxicity signs and motor activity (spaces roamed [S]). Reflexes (grooming or piloerection) were assessed after 24 h. Mortality, body weight, and water and food intake were concurrently monitored for 14 days. At the end of this period, animals were euthanized in a CO₂ chamber, and the heart, lung, liver, and kidney of animals were dissected and weighed (normalized to body weight⁹), and processed for hematoxylin-eosin histological analysis. Histopathological images were obtained using a camera-equipped microscope and analyzed with NS-Elements D software. This study was approved by the ethics committee on animal use of Universidade Nove de Julho (opinion # AN 0003/11).

Fasting (18 h) Wistar rats (250 - 350g) were decapitated, and the ileum was carefully isolated and dissected⁸. Tissue strips (1.5 cm) were mounted in organ baths containing 5 mL of modified Krebs-Henseleit solution at 37 °C with 11.0 mM glucose, and continuously aerated with O₂¹⁰. Strips were connected to a force transducer and amplifier, and contraction data were recorded using AQCD software. A Cc-EtOH stock solution (10 mg/mL) was prepared in 0.1% cremophor and diluted in Mili-Q water. After 30 min of equilibration period, strips were contracted with 1 μM carbachol (CCh) or 40 mM KCl, following 15 min of incubation with Cc-EtOH (27-730 μg/mL) or vehicle. Contractile responses to CCh or KCl were compared in the absence and presence of Cc-EtOH extract. The IC₅₀ of the Cc-EtOH extract in CCh- or KCl-induced contraction were calculated by non-linear regression¹¹. This protocol was approved by the ethics committee on animal use of the Universidade Federal de São Paulo (opinion number 4295060514/14).

Results were expressed as mean ± SEM. Student T-test or one-way ANOVA test followed by post-hoc Dunnett test for multiple comparisons were used when applicable. Differences were considered significant at p < 0.05.

RESULTS

Oral administration of ethanol induced stomach lesions (UI = 368 ± 89 mm²; Figure 1). Pretreat-

ment with 50 mg/kg Cc-EtOH extract (UI = 210.0 ± 55.8 mm²) did not prevent the ethanol-induced lesions compared with the control group. However, treatments with 150 and 500 mg/kg Cc-EtOH pre-

vented ethanol-induced lesions in a dose-dependent manner (UI = 119 ± 39 and 48 ± 13 mm², respectively), similar to the effect observed with 20 mg/kg omeprazole (UI = 146 ± 20 mm²).




Figure 1. Ethanol-induced lesions (control) in the presence of omeprazole (20 mg/kg) or Cc-EtOH extract (50, 150, and 500 mg/kg) on rat stomachs (n = 6).

Histological analysis confirmed epithelial damage and inflammatory infiltration in the control group (Figure 2A), consistent with the characteristic mucosal injury in the ethanol-induced ulcer lesion model. However, the single-dose treatment with 500 mg/kg Cc-EtOH extract prevented ethanol-induced gastric lesions (Figure 2B). Pretreatment also attenuated the inflammatory response, suggesting regeneration and re-epithelization of the gastric mucosa.

In subsequent experiments, oral administration of 2 g/kg Cc-EtOH extract did not induce mortality or changed body weight, water or food intake (data not shown). However, treated animals showed significantly reduced ambulation between 30 and 60 min after treatment ($S = 47.8 \pm 4.4$ and 34.4 ± 3.1 units, respectively) compared with the control group (Figure 3). Normalized organ weights are presented in Table 1.





Figure 2. Effects of Cc-EtOH on ethanol-induced rat gastric lesions stained with hematoxylin and eosin. Panel A shows a photomicrograph (100 x) of epithelial lesion and inflammatory cell infiltration in the gastric mucosa after oral administration of ethanol. Panel B shows the effect of pretreatment with 500 mg/kg Cc-EtOH extract, preventing the instauration of epithelial lesions or inflammatory infiltration in the gastric mucosa.

Table 1. Weight of the indicated organs isolated from mice treated or not (control) with Cc-EtOH extract.

Treatment	Weight \pm SEM (mg/g)			
	Liver	Heart	Lung	Kidney
Control	7.7 \pm 0.7	0.9 \pm 0.1	1.9 \pm 0.1	2.2 \pm 0.1
Cc-EtOH	6.5 \pm 0.3	0.6 \pm 0.1*	1.0 \pm 0.2**	2.0 \pm 0.1


Data are presented as mean \pm SEM. (n = 5, t-test *p < 0.05; **p < 0.01)

**Figure 3.** Effect of Cc-EtOH extract on mice ambulation from 30 min to 120 min of Cc-EtOH treatments (n = 5).

No significant changes were observed in liver or kidney weights, whereas heart and lung weights were reduced in treated animals compared with control group.

Preincubation with Cc-EtOH extract equipo-

tently antagonized the CCh- and KCl-induced phasic contractions (Figure 4A and B, respectively). No significant differences were observed between the calculated IC₅₀ for the Cc-EtOH extract in CCh-induced contraction (236 \pm 59 μ g/mL) and KCl-induced contractions (201 \pm 43 μ g/mL).

**Figure 4.** Effects of Cc-EtOH extract on phasic contractions induced by 1 μ M CCh (panel A) or 40 mM KCl (panel B) of rat isolated ileum (n = 4).

DISCUSSION

The Cc-EtOH extract exhibited anti-ulcer activity in rats, showed no acute toxicity in mice, and presented antispasmodic effects in isolated rat ileum *in vitro*.

Considering the traditional use of *C. corindum* (*balãozinho*) in Bahia, Brazil, for its anti-ulcerative properties, we investigated its activity using an

ethanol-induced stomach ulcer model in rats⁸. The Cc-EtOH extract protected the gastric mucosa from ethanol-induced lesions with efficacy comparable to omeprazole, a proton pump inhibitor (Figure 1). This protective effect is consistent with previous findings in *C. halicacabum*¹¹ and *Serjania caracasana*, two other *Sapindaceae* species¹³. Thus, the anti-ulcer properties of *C. corindum* may be associated with

its chemical composition, including triterpenes, flavonoids, and steroids^{4,5}, also reported in related species¹³.

No toxicological data were previously available for *C. corundum*, underscoring the need for *in vivo* assays. In the present study, acute oral toxicity of Cc-EtOH extract was investigated in mice, an appropriate model for initial toxicity screening¹⁴. No toxicity was detected, as no mortality or tissue damage was observed during the 14-day evaluation period. However, transient sedative effects were evidenced by decreased locomotor activity (Figure 3), consistent with earlier reports of central nervous system effects of this plant¹⁵. Nevertheless, this reduction in ambulation was limited to the first 60 min following oral administration. Organ weight analysis revealed decreased heart and lung weights (Table 1), although no histological alterations were observed. While the physiological relevance of these evidence remains unclear, the absence of morphological changes supports the safety of a single 2 g/kg oral dose of Cc-EtOH extract. Notably, this dose was four-fold higher than the effective anti-ulcer dose (500 mg/kg; Figure 1) and did elicit putative toxicity.

Further investigation in the gastrointestinal tract demonstrated that Cc-EtOH extract inhibited CCh- or KCl-induced phasic contractions in isolated ileum strips *in vitro* with similar potency, suggesting effects on two distinct transduction pathways involved in intestinal contraction. These contractile responses, whether pharmaco-mechanical or electromechanical couplings rely on Ca²⁺ influx through voltage-dependent Ca²⁺-channels¹⁶.

The observed *in vivo* gastroprotective may, at least in part, rely on the antispasmodic proprieties of Cc-EtOH extract *in vitro*, similar to agents such as atropine (i.e., reducing gastrointestinal motility) and prostaglandins (i.e., protecting gastric mucosa)¹⁷.

In conclusion, Cc-EtOH extract promoted gastric protection in rats and antispasmodic effects *in vitro*, without evidence of acute toxicity. Together, these findings strongly support the traditional use of *balãozinho* in folk medicine for the treatment of gastric ulcers.

CONFLICTS OF INTEREST

Nothing to declare.

REFERENCES

- Jainu M, Devi CSS. Antiulcerogenic and ulcer healing effects of *Solanum nigrum* (L.) on experimental ulcer models: Possible mechanism for the inhibition of acid formation. *J Ethnopharmacol.* 2006; 104:56–163.
- Somner GV, Carvalho ALG, Siqueira CT. Sapindaceae da Restinga da Marambaia, Rio de Janeiro, Brasil. *Rodrigueusia.* 2009; 60: 485-507.
- Agra MF, KN Silva, Basílio Ionaldo JLDB, Freitas PF, Barbosa-Filho JM. Survey of medicinal plants used in the region Northeast of Brazil. *Rev Bras Farmacogn.* 2008; 18(3): 472-508.
- Adinarayana D, Sarada M, Ramachandraiah P. Occurrence of a rare flavone glucuronide in *Cardiospermum corindum*. *Indian J Nat Prod.* 1989; 5(2): 8-10.
- Silva FL, Moreno PRH, Braz-Filho R, Tavares JF, Barbosa-Filho JM. Chemical constituents of *Cardiospermum corindum* L. and their distribution in Sapindaceae. *Biochem Syst Ecol.* 2014; 57: 137-40.
- Morimoto Y, Shimohara K, Oshima S, Sukamoto T. Effects of the new anti-ulcer agent KB-5492 on experimental gastric mucosal lesions and gastric mucosal defensive factors, as compared to those of terpenone and cimetidine. *Jpn J Pharmacol.* 1991; 57(4):595-605.
- Kauffman GL, Grossman MI. Prostaglandin and cimetidine inhibit the formation of ulcers produced by parenteral salicylates. *Gastroenterology.* 1978; 75: 1099-1102.
- Walker R, Wilson KA. Prostaglandins and the contractile action of bradykinin on the longitudinal muscle of rat isolated ileum. *Br J Pharmacol.* 1979; 67: 527-33.
- Sun YD, Benishin CG. K⁺ channel openers relax longitudinal muscle of guinea-pig ileum. *Eur J Pharmacol.* 1994; 271: 453-59.
- Neubig RR, Spedding M, Kenakin T, Christopoulos A. International union of pharmacology committee on receptor nomenclature and drug classification. IX. Recommendations on terms and symbols in quantitative pharmacology. *Pharm Rev.* 1995;47(2):255-66.
- Sheeba MS, Asha VV. Effect of *Cardiospermum halicababum* on ethanol-induced gastric ulcers in rats. *J Ethnopharmacol.* 2006; 106: 105–10.
- Silva JLV, Carvalho VS, Silva FL, Barbosa-Filho JM, Rigoni VLS, Nouailhetas VLA. Gastrointestinal property of *Serjania caracasana* (Jacq) Willd. (Sapindaceae) on rats. *Pharmacol On Line.* 2012; 1: 22-6.
- Heredia-Vieira SC, Simonet AM, Vilegas W, Macías FA. Unusual C, O-Fused Glycosylapigenins from *Serjania marginata* Leaves. *J Nat Prod.* 2014; 78(1):77-84.
- The Organisation Of Economic Co- Operation Development (OECD). OECD guideline for testing of chemicals. 420 Acute Oral Toxicology – Fixed Dose Procedure. The Organisation of Economic Co-operation Development (OECD), 200: 1-14.
- Bhattacharya SK, Satyan KS. Experimental methods for evaluation of psychotropic agents in rodents: I-Anti- anxiety agents. *Indian J Exp Biol.* 1997; 35(6): 565-75.
- Sanders KM. Regulation of smooth muscle excitation and contraction. *Neurogastroenterol Motil.* 2008; 20(1):39-53.
- Ukawa H, Sugawa Y, Yagi K, Suzuki K, Takeuchi K. The

ORIGINAL ARTICLE

roles of prostaglandin E receptor subtypes in the cytoprotective action of prostaglandin E2 in rats stomach Aliment Pharmacol Ther. 2000;14(1):116-24.